

On the parameters of the QCD-motivated potential in the relativistic independent quark model

V.V. Khrushev^{a,c}, V.I. Savrin^b and S.V. Semenov^{c, 1}

^aCenter for Gravitation and Fundamental Metrology, VNIIMS, Moscow, Russia

^bInstitute of Nuclear Physics, Moscow State University, Russia

^cRussian Research Center "Kurchatov Institute", Kurshatov Square 1, Moscow, 123182, Russia

Abstract

In the framework of the relativistic independent quark model the parameters of the QCD-motivated static potential and the quark masses are calculated on the basis of the 1^{--} meson mass spectra. The value of the confining potential coefficient is found to be $(0.197 \pm 0.005) \text{ GeV}^2$ for quark-antiquark interaction independently on their flavours. The dependence of the quasi-Coulombic potential strength on the interaction distance are consistent with the QCD-motivated behaviour. The $q\bar{q}$ -separations are evaluated and the e^+e^- decay widths are estimated with the help of relativistic modification of the Van Royen-Weisskopf formula.

PACS: 12.39.Pn; 12.39.Ki; 12.40.Yx

Keywords: Quark model; Potential; Meson masses; Decay widths; Strong coupling constant; String tension; Quark masses; Radial excitations

¹E-mail: semenov@imp.kiae.ru

The calculation of spectroscopical hadron characteristics within the data precision [1] still remains among the unsolved QCD problems. The main difficulty consists in consideration of the nonperturbative QCD effects, which are not understood yet from first principles but only in terms of various QCD-inspired models. Thus, at the present time phenomenological hadron models can be used to a certain extent for the data interpretation. Among phenomenological hadron models of different kinds (e.g., Refs. [2-8]) the relativistic model for quasi-independent quarks seems to be one of the most interesting and simplest [9]. In our previous papers [10–13] we developed the translation invariant version of this model for description of spectroscopical characteristics of both the light and heavy vector mesons. We used the Dirac equation with a static QCD-motivated potential to describe the quark or antiquark motion in the mean field inside the meson. The model does not contradict the first principles of QCD and allows one in a simple fashion to carry out numerical calculations of the meson characteristics, such as their masses, the average separations between quarks and antiquarks and the e^+e^- decay widths of vector mesons. However, in those papers the effective potential entered in the second order model radial equation was chosen for simplicity in an approximate form. In particular this simplification lead to the constancy of the quasi-Coulombic potential strength evaluated on the basis of 1^{--} meson mass spectra for different quark flavours.

In the present paper in the framework of the relativistic independent quark model [10–13] we use the exact expression for the static interaction effective potential and evaluate the potential parameters coupled with the quark masses on the basis of 1^{--} meson mass spectra. Then the obtained values are used for calculating the masses of 1^{--} radial excitations. The e^+e^- decay widths of heavy mesons are estimated with the help of relativistic modification of the Van Royen-Weisskopf formula. The $q\bar{q}$ -separations for the light and heavy mesons are evaluated as well. We find an alternation of the strong coupling constant α_s (which depends on the quark flavour and is consistent with the QCD asymptotic freedom phenomenon) together with the constancy of the confining potential coefficient for all quark flavours. The latter result supports the hypothesis of the flavor independence of confinement potential [14–16].

According to the main statement of the independent quark model the hadron is considered as a system composed of the non-interacting with each other directly valent constituents (quarks and antiquarks) moving in some mean field. One supposes that this field is a colour singlet confining field, which is produced by the constituents and takes into account the effects of creation and annihilation of the sea $q\bar{q}$ pairs as well. To simplify calculations it is assumed that this mean field is spherically symmetric and its motion in space is determined by motion of its center. Furthermore, in order to keep out the ghost-motion states of the center of mass for the independent constituents, we associate the center of mass with the mean field center. We treat the mean field as a quasi-classical object possessing some energy, however one cannot evaluate the mean field energy without additional assumptions, and this quantity is a phenomenological parameter in the model considered. Each of constituents interacting with the spherical mean field gets the state with a definite value of its energy. On the phenomenological ground the mass formula for J^{PC} meson, which is $n^{2S+1}L_J$ state of $q\bar{q}'$ -system, can be presented in the following form [10–13]:

$$M_n(J^{PC}) = E(n_r, j) + E(n'_r, j'), \quad (1)$$

where $n_r(n'_r)$ and $j(j')$ are the radial quantum number and the angular momentum quantum number of the quark (antiquark). $E(n_r, j)$ is called the energy spectral function or the mass term for the quark and in the framework of our model it has got the following phenomenological expression:

$$E(n_r, j) = [\lambda + m^2]^{1/2} + c[1 + (-1)^{L+j+1/2}], \quad (2)$$

The first term represents the relativistic effective energy of the constituent moving in the mean field inside the meson, the quark mass m being a model parameter and λ being found as an eigenvalue of the radial equation of the model. The second term in the formula (2) includes a part of energy of the mean field and is purely phenomenological one, which gives small corrections to the spectra of different meson families.

In the equal-time approximation the entire system, the vector $q\bar{q}$ meson, is described by a stationary wave function in the center-of-mass frame, which in the translation-invariant model [10–13] can be represented as follows:

$$\Phi_\eta(\mathbf{r}, \mathbf{r}') = \phi_0 \bar{\psi}(\mathbf{r}) e_\eta^\mu \gamma_\mu U_C \bar{\psi}(\mathbf{r}'), \quad (3)$$

where $\mathbf{r}$ and $\mathbf{r}'$ are the quark and antiquark coordinates in respect to the mean field center, ϕ_0 is some constant, e_η is a 4-vector of the meson polarization, η describes a meson polarization state and takes the values (1, 0, -1), U_C is the charge conjugation operator. Thus, a constituent turns out to be coupled to the other via the common mean field which is moving together with their center of mass.

The wave function $\psi(\mathbf{r})$ for any constituent (the quark or antiquark orbital) is a solution of the single-particle equation with a static interaction potential. We choose the Dirac equation with the QCD-motivated potential $V(r)$ for description of interaction of the quark or antiquark with the external mean field in order to determine the orbitals inside the meson. The potential $V(r)$ is spherically symmetrical and consists of the Lorentz scalar and vector parts: $V(r) = \beta V_0(r) + V_1(r)$. Hence the equation for a fermionic constituent is

$$\sqrt{\lambda + m^2} \psi(\mathbf{r}) = [(\boldsymbol{\alpha} \mathbf{p}) + \beta(m + V_0) + V_1] \psi(\mathbf{r}) \quad (4)$$

with $V_0(r) = \sigma r/2$ and $V_1(r) = -2\alpha_s/3r$, where the two model parameters σ and α_s are introduced and have got meanings of the string tension and the strong coupling constant at small distances, correspondingly.

It is well-known that the solutions of Eq. (4) with the total angular momentum j and its projection m can be represented as

$$\psi(\mathbf{r}) \propto \begin{pmatrix} f(r) \Omega_{jl}^m(\mathbf{n}) \\ -ig(r) (\boldsymbol{\sigma} \mathbf{n}) \Omega_{jl}^m(\mathbf{n}) \end{pmatrix}, \quad (5)$$

where $\mathbf{n} = \mathbf{r}/r$. If $k = -\omega(j + 1/2)$ where ω is an eigenvalue of the space-parity operator, the system of the radial Dirac equations for the fermion in the mean field with the definite energy sign and spin projection reads

$$\begin{aligned}
(\sqrt{\lambda + m^2} - V_0 - V_1 - m)f &= -\frac{(k+1)}{r}g - g', \\
(\sqrt{\lambda + m^2} + V_0 - V_1 + m)g &= -\frac{(k-1)}{r}f + f'.
\end{aligned} \tag{6}$$

Using Eqs. (6) one can derive the second order equation for the "large" component $f(r)$, and then making a substitution

$$\varphi(r) = rf(r) \left[V_0(r) - V_1(r) + m + \sqrt{\lambda + m^2} \right]^{-1/2} \tag{7}$$

one comes on to the model radial equation for $\varphi(r)$ in the following form:

$$\begin{aligned}
\varphi'' + \lambda\varphi &= \left[(m + V_0)^2 - (\sqrt{\lambda + m^2} - V_1)^2 + \frac{k(k-1)}{r^2} + \frac{3(V_0' - V_1')^2}{4(\sqrt{\lambda + m^2} - V_1 + V_0 + m)^2} \right. \\
&\quad \left. - \frac{k(V_0' - V_1')}{r(\sqrt{\lambda + m^2} - V_1 + V_0 + m)} - \frac{(V_0'' - V_1'')}{2(\sqrt{\lambda + m^2} - V_1 + V_0 + m)} \right] \varphi.
\end{aligned} \tag{8}$$

Further on we restrict ourselves evaluating only characteristics of the radial excitations of the S-wave 1^{--} mesons because they are described by the simplest version of model and supported by the most extensive set of accurate data [1], especially for heavy mesons. So, $k = 1$ and the contribution of the term containing c in Eq. (2) is equal to zero. λ entering Eq. (2) can be calculated with the help of the S-wave radial model equation:

$$\begin{aligned}
\varphi'' + \lambda\varphi &= \left\{ -\frac{4\alpha_s\sqrt{\lambda + m^2}}{3r} - \left(\frac{2\alpha_s}{3r} \right)^2 + m\sigma r \right. \\
&\quad \left. + \left(\frac{\sigma r}{2} \right)^2 - \left[\frac{\sigma r}{2} (m + \sqrt{\lambda + m^2}) + \left(\frac{\sigma r}{4} \right)^2 + \frac{5\alpha_s\sigma}{6} - \frac{\alpha_s^2}{3r^2} \right] \right. \\
&\quad \left. \left[\frac{2\alpha_s}{3} + r (m + \sqrt{\lambda + m^2}) + \frac{\sigma r^2}{2} \right]^2 \right\} \varphi.
\end{aligned} \tag{9}$$

The right-hand side of Eq. (9) has a singularity at the origin, and when $r \rightarrow 0$ it behaves as $3/4r^2 - 4\alpha_s^2/9r^2$. Therefore one should keep $\alpha_s < 3/2$ in order to prevent a fall at the origin. At the infinity $r \rightarrow \infty$ the effective potential behaves as the oscillatory one and tends to $\sigma^2 r^2/4$.

In the framework of our model the quantities m , σ and α_s are phenomenological parameters and to be determined as a result of calculations of meson mass spectra, as well as the e^+e^- decay widths, and a comparison of them with experimental data.

The model equation (9) can be solved only by numerical methods. For calculating its eigenvalues we used the computer code algorithm, which is based on the Numerov three point recurrent relation for the equation $y'' = F(r, E)y$ [17]:

$$y_{i-1} = \left[y_i \left(2 + \frac{5}{6} F(r_i, E) h^2 \right) - y_{i+1} \left(1 - \frac{F(r_{i+1}, E) h^2}{12} \right) \right] \times \left[1 - \frac{F(r_{i-1}, E) h^2}{12} \right]^{-1}. \quad (10)$$

For bound states $y(r)$ tends to zero when $r > r_{cl}$. Here r_{cl} is a classical radius of the bound state, which is determined from the equation $U(r_{cl}) = E$ and E is the initial energy of the considered level. Then the E value can be determined from the condition: $y(0) = 0$. However, $F(r, E)$ has a singularity at the origin. So, when we calculate $y(0)$ with the help of formula (10) we use a regularization procedure for the singular potential, which is analogous to the procedure worked out, for instance, in Ref. [18].

In order to estimate the precision of the calculation algorithm the well-known test with diminished spacing h is used. The results obtained allow us to disregard machine calculation errors as compared with systematic errors of the model. The accuracy criterion for fitted parameters was the maximum value of the acceptable errors for evaluated hadron mass values as compared with typical experimental errors which we choose to be in the range of $30 \div 40$ MeV. Thus, the parameter errors written below must be considered as our estimation of systematic errors of the model.

When calculating the values of the model parameters we do not suppose *a priori* the validity of the hypothesis of flavor independence for the confining potential. First of all the model parameters m , α_s and σ for the $b\bar{b}$ and $c\bar{c}$ radial excitations of 1^{--} states were found. The fit was carried out for each meson family independently. The existing experimental data allowed to find these parameters as well as to prove the validity of the model. In this manner it was found that the value of σ is the same for the $b\bar{b}$ and $c\bar{c}$ states within the systematic errors of the model and equal to (0.197 ± 0.005) GeV². Taking into account that there are no well established data for the higher radial excitations of 1^{--} light mesons, the obtained σ value was used for calculating m and α_s for light quarks. Note that when doing the calculations we neglect the isotopic mass splitting of the u - and d -quarks because it is beyond of the model accuracy.

All results for the well known experimental data on the 1^{--} mesons, which are composed of the quark and antiquark with u -, d -, s -, c - or b -flavours, are consistent within the model accuracy with the following values for the coupling α_s (quasi-Coulombic potential strength) and the quark masses:

$$\begin{aligned} \bar{m}_{u,d} &= (0.01 \pm 0.008) \text{ GeV}, & \alpha_s^{u,d} &= 0.70 \pm 0.15, \\ m_s &= (0.18 \pm 0.03) \text{ GeV}, & \alpha_s^s &= 0.50 \pm 0.10, \\ m_c &= (1.35 \pm 0.05) \text{ GeV}, & \alpha_s^c &= 0.36 \pm 0.03, \\ m_b &= (4.70 \pm 0.10) \text{ GeV}, & \alpha_s^b &= 0.24 \pm 0.02. \end{aligned}$$

Now let us estimate the e^+e^- decay widths of the vector mesons with the help of solutions of the Dirac equations (6) which were obtained numerically. It is well known that in the nonrelativistic potential approach the vector meson e^+e^- width is described by the Van Royen-Weisskopf formula with first order QCD correction:

$$\Gamma_V = \frac{16\pi\alpha^2 e_q^2}{M_V^2} |\Psi(0)|^2 \left(1 - \frac{16\alpha_s}{3\pi} \right), \quad (11)$$

where $|\Psi(0)|^2$ is the square of the meson wave function, which describes the probability density of meson at the origin. We shall keep a structure of the formula (11) in the relativistic independent quark model.

Using Eqs. (3) and (5) the wave function of the meson consisting of independent quark and antiquark in the S -state can be written as

$$\Phi_\eta(\mathbf{r}, \mathbf{r}') = \frac{\phi_0}{4\pi} \left[-f(r)f(r') \overset{*}{w}(\boldsymbol{\sigma}\mathbf{e}_\eta)w' + g(r)g(r') \overset{*}{w}(\boldsymbol{\sigma}\mathbf{n})(\boldsymbol{\sigma}\mathbf{e}_\eta)(\boldsymbol{\sigma}\mathbf{n}')w' \right], \quad (12)$$

where we have used a relationship $\sigma_2 \overset{*}{\boldsymbol{\sigma}} = -\boldsymbol{\sigma}\sigma_2$ and denoted $w' = -\sigma_2 \overset{*}{w}$ (the asterisk denotes complex conjugation). So the probability density of independent quark-antiquark system can be easily calculated:

$$\begin{aligned} |\Phi_\eta(\mathbf{r}, \mathbf{r}')|^2 &= \frac{2\phi_0^2}{16\pi^2} \{ f^2(r)f^2(r') + g^2(r)g^2(r') + 2f(r)g(r)f(r')g(r')(\mathbf{n}\mathbf{n}') - \\ &\quad - 2f(r)g(r)f(r')g(r')[(\mathbf{e}_\eta\mathbf{n})(\overset{*}{\mathbf{e}}_\eta\mathbf{n}') + (\overset{*}{\mathbf{e}}_\eta\mathbf{n})(\mathbf{e}_\eta\mathbf{n}')]\}. \end{aligned} \quad (13)$$

And the averaged over initial spin projections probability density is

$$|\overline{\Phi_\eta(\mathbf{r}, \mathbf{r}')}|^2 = \frac{2\phi_0^2}{16\pi^2} [f^2(r)f^2(r') + g^2(r)g^2(r') + 2/3 f(r)g(r)f(r')g(r')(\mathbf{n}\mathbf{n}')]. \quad (14)$$

Now taking into account that $\mathbf{r}$ and $\mathbf{r}'$ are the coordinates of the quark and the antiquark in respect to their center of mass, which coincides with the center of the mean field, we can define according to Eq. (14) the probability density of meson in the following way:

$$\int d^3\mathbf{r}' |\overline{\Phi_\eta(\mathbf{r}, \mathbf{r}')}|^2 = \frac{1}{2\pi} [\kappa_1 f^2(r) + \kappa_2 g^2(r)], \quad (15)$$

where we denote

$$\int dr r^2 f^2(r) = \kappa_1, \quad \int dr r^2 g^2(r) = \kappa_2. \quad (16)$$

Hence the full probability is

$$\int d^3\mathbf{r} \int d^3\mathbf{r}' |\overline{\Phi_\eta(\mathbf{r}, \mathbf{r}')}|^2 = 2(\kappa_1^2 + \kappa_2^2). \quad (17)$$

and therefore the normalized probability density which is analogous to the $|\Psi(r)|^2$ is equal to $(\kappa_1 f^2(r) + \kappa_2 g^2(r)) / 4\pi(\kappa_1^2 + \kappa_2^2)$. This quantity at $r = 0$ must enter the formula (11) instead of the $|\Psi(0)|^2$. However, in relativistic theory one cannot localized a quark within the volume with the radius less than the Compton wave length of the quark $\lambda_q \cong m_q^{-1}$. According to this reasoning we use for the estimation of decay width the following formula:

$$\Gamma_{V_n} = \frac{4\alpha^2 e_q^2}{M_{V_n}^2} \left[\frac{\kappa_1 f_n^2(\lambda_q) + \kappa_2 g_n^2(\lambda_q)}{\kappa_1^2 + \kappa_2^2} \right] \left(1 - \frac{16\alpha_s}{3\pi} \right). \quad (18)$$

Taking into account the expression for the normalized probability density, the mean squared radius of the meson is

$$R_n^2 = \int dr r^4 \left[\frac{\kappa_1 f_n^2(r) + \kappa_2 g_n^2(r)}{\kappa_1^2 + \kappa_2^2} \right]. \quad (19)$$

The results obtained for the meson masses M_n^{th} and the average radii R_n^{th} are displayed in Table 1. We estimate the e^+e^- decay widths Γ_n^{th} with the help of formula (18) only for the $c\bar{c}$ - and $b\bar{b}$ -mesons because there are no satisfactory data for the radial excitations of light mesons. The mass values M_n^{th} and M_n^{exp} coincide with each other within the model accuracy. Taking into account the approximate nature of the formula (18) there is a conformity between the calculated decay ratios and the experimental values [1]. However, some further efforts are needed in order to improve the precision for decay widths calculations in the framework of model considered. Note that there is a good agreement between the R_n^{th} and the values calculated in the framework of the lattice QCD in the valence quark approximation [19].

Table 1

In this paper we considered the relativistic hadron model with independent quarks and evaluated the parameters of the $q\bar{q}$ model potential, as well as the masses, the average radii of the 1^{--} mesons, and estimated the e^+e^- decay widths. We find that the value of $\sigma = (0.197 \pm 0.005) \text{ GeV}^2$ is the admissible string tension value for u -, d -, s -, c - and b -quark flavours within the systematical errors of the model. It is an important result of our paper which confirms the flavor independence of confining potential on the $3 \cdot 10^{-2}$ level of accuracy. Thus one may interpret σ as the universal parameter of quark-antiquark interaction at large distances and consider the characteristic confinement scales such as the confinement length l_c and the confinement mass m_c , which satisfy the following relations: $m_c l_c = 1$, $\sigma = m_c^2$.

In addition, the correspondence to the QCD asymptotic freedom phenomenon, or the decrease of α_s when the interaction range diminishes, is confirmed both for the light and heavy quarks in the framework of this model. Note that within the large systematical errors of the model it is difficult to prove the logarithmic dependence of α_s on Q^2 , moreover, there are considerable nonperturbative contributions for the light quarks (see, e.g. Refs. [20, 21]). Nevertheless, there is no contradiction between the $\alpha_s(Q^2)$ behaviour in QCD and the diminution of the parameter α_s in this model when coming from the heavy to the light quarks.

An advantage of the presented model is its conceptual clarity and technical simplicity, which allows one to calculate in a simple fashion the meson characteristics such as masses, $q\bar{q}$ -separations and e^-e^+ decay widths with an admissible among phenomenological models accuracy. For instance, the values of the parameters m_b and m_c evaluated in the framework of this model are similar to the values of the pole masses M_b and M_c obtained in Ref. [7]. The values of $\alpha_s(m_i^2)$ ($i = b, c$) are close to the $\alpha_s(M_i^2)$ in the $\overline{MS}$ scheme, whereas the average $q\bar{q}$ -separations for heavy vector mesons are in agreement with those found on the basis of the lattice QCD [19]. Moreover, in this paper the calculations of $q\bar{q}$ -separations for light vector mesons are presented, while the same calculations are not possible now in the frame of the lattice QCD. In order to increase the model accuracy it is reasonable to take into account the relativistic corrections due to contact particle-particle interaction, like those which have been treated in Ref. [2]. The account for such kind of corrections lies beyond the scope of the independent quark model and will be of our further task in

order to improve the model accuracy and to find its limitations. Besides we intend to apply this model to the more complicated systems, such as the baryons and the exotic multiquark mesons.

Acknowledgments. The authors express their gratitude to R.N. Faustov and A.L. Kataev for useful discussions.

References

- [1] D.E. Groom et al. (Particle Data Group), The European Phys. Journ. C15 (2000) 1.
- [2] S. Godfrey and N. Isgur, Phys.Rev., D 32 (1985) 189.
- [3] W. Lucha, F.F. Schoberl and D. Gromes, Phys.Rep., 200 (1991) 128.
- [4] S. Titard and F.J. Yndurain, Phys.Rev., D51 (1995) 6348.
- [5] S.G. Gorishny, A.L. Kataev and S.A. Larin, Phys. Lett. B 135 (1984) 458.
- [6] A. Duncan, E.Eichten and H. Thacker, Phys. Lett. B 303 (1993) 109.
- [7] M. Jamin and A. Pich, Nucl. Phys. B507 (1997) 334.
- [8] G. Jaczko and L. Durand, Phys. Rev. D 58 (1998) 114017.
- [9] P.N. Bogoliubov and A.E. Dorokhov, Part. & Nucl. 8 (1987) 917.
- [10] V.V. Khrushev, Yad. Fiz. 46 (1987) 219; *ibid.* 55 (1992) 773.
- [11] V.I. Savrin, S.V. Semenov and V.V. Khrushev, in: Problems on High Energy Physics and Field Theory (Nauka, Moscow, 1989) p. 224.
- [12] S.V. Semenov and V.V. Khrushev, Yad. Fiz. 56 (1993) 218.
- [13] V.V. Khrushev, V.I. Savrin and S.V. Semenov, Phys. Lett. B374 (1996) 159.
- [14] A. E. Dorokhov, JINR preprint R2-121159, Dubna (1979).
- [15] A. Martin, CERN preprint TH-2741, Geneva (1979).
- [16] C. Quigg, J.L. Rosner and H.B.Thacker, FERMILAB preprint 79-52 (THY), Batavia (1979).
- [17] B.V. Numerov, Proc. Centr. Astrophys. Observ. 2 (1933) 188.
- [18] E. Buendia and R. Guardiola, J. Comput. Phys. 60 (1985) 561.
- [19] G.S. Bali, K. Schilling and A. Wachter, Phys. Rev. D56 (1998) 2566.
- [20] D.V. Shirkov and I.L. Solovtsov, Phys. Rev. Lett. 79 (1997) 1209.
- [21] A.I. Alekseev and B.A. Arbuzov, Mod. Phys. Lett. A 13 (1998) 1747.

Table 1. Evaluated masses and e^+e^- decay widths of the 1^{--} mesons in comparison with the data from Ref. [1], and the average $q\bar{q}$ -separations in comparison with the results of Ref. [19].

Meson	$M_n^{exp}[MeV]$	$M_n^{th}[MeV]$	$R_n^{th}[fm]$	$R_n^{[19]}[fm]$	$\Gamma_n^{th}[keV]$	$\Gamma_n^{exp}[keV]$
ρ	769.9 ± 0.8	740	0.65	-	-	6.77 ± 0.32
ρ'	1465 ± 25	1455	1.12	-	-	-
ρ''	1700 ± 20	1730	1.32	-	-	-
ϕ	1019.413 ± 0.008	1010	0.58	-	-	4.43 ± 0.05
ϕ'	1680 ± 50	1650	0.96	-	-	-
ϕ''	-	2050	1.34	-	-	-
J/ψ	3096.88 ± 0.04	3060	0.41	0.43	7.32	5.26 ± 0.37
ψ'	3686.00 ± 0.09	3650	0.79	0.85	2.03	2.14 ± 0.21
ψ''	4040 ± 10	4070	1.06	1.18	0.97	0.75 ± 0.15
ψ'''	4415 ± 6	4390	1.30	1.47	0.58	0.47 ± 0.10
Υ	9460.37 ± 0.21	9470	0.26	0.24	1.30	1.32 ± 0.05
Υ'	10023.3 ± 0.3	9990	0.54	0.51	0.37	0.520 ± 0.032
Υ''	10355.3 ± 0.5	10325	0.75	0.73	0.21	-
Υ'''	10580 ± 3.5	10550	0.94	0.93	0.16	0.248 ± 0.031
Υ^{5S}	10865 ± 8	10830	1.10	-	0.12	0.31 ± 0.07
Υ^{6S}	11019 ± 8	10985	1.25	-	0.08	0.13 ± 0.03